



Generalizations of a parameterized Jordan-type inequality, Janous's inequality and Tsintsifas's inequality

Shanhe Wu^a, Lokenath Debnath^{b,*}

^a Department of Mathematics, Longyan College, Longyan Fujian 364012, People's Republic of China

^b Department of Mathematics, University of Texas-Pan American, Edinburg, TX 78539, USA

ARTICLE INFO

Article history:

Received 22 January 2008

Accepted 22 January 2008

Keywords:

Jordan's inequality

Janous's inequality

Tsintsifas's inequality

Parameterized Jordan-type inequality

Sharpening

Generalization

ABSTRACT

In this work, we first prove a generalized version of a parameterized Jordan-type inequality. We then use it to prove the generalized versions of Janous's inequality and Tsintsifas's inequality which reduce to two inequalities conjectured by Janous and Tsintsifas as special cases.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

The celebrated Jordan's inequality is stated as [Theorem A](#) (see [Mitrinović and Vasić \[1, p.33\]](#)).

Theorem A. Let $0 < x \leq \frac{\pi}{2}$. Then

$$\frac{2}{\pi} \leq \frac{\sin x}{x} < 1. \quad (1)$$

Jordan's inequality (1) and its modified versions have many important applications in calculus, trigonometry and the theory of limits. In recent years, this classical inequality has received considerable attention, different proofs, various generalizations, improvements and analogues have been presented by many authors (see [\[2–17\]](#) and references therein).

Recently, Janous [\[18\]](#) and Tsintsifas [\[19\]](#) proposed the following inequalities involving the ratio $(\sin x)/x$ as conjectures:

Conjecture 1 (See Janous [\[18\]](#)). Suppose $A > 0$, $B > 0$, $C > 0$ and $A + B + C = \pi$. Then the following inequality holds true:

$$2 < \frac{\sin A}{A} + \frac{\sin B}{B} + \frac{\sin C}{C} \leq \frac{9\sqrt{3}}{2\pi}. \quad (2)$$

Conjecture 2 (See Tsintsifas [\[19\]](#)). Suppose $0 < A < \pi/2$, $0 < B < \pi/2$, $0 < C < \pi/2$ and $A + B + C = \pi$. Then the following inequality holds true:

$$\frac{3}{\pi} < \frac{\sin A}{\pi - A} + \frac{\sin B}{\pi - B} + \frac{\sin C}{\pi - C} < \frac{3\sqrt{3}}{\pi}. \quad (3)$$

* Corresponding author.

E-mail addresses: wushanhe@yahoo.com.cn (S. Wu), debnathl@utpa.edu (L. Debnath).

Recently, Klamkin [20] and Murty et al. [21] have given affirmative proofs of [Conjectures 1](#) and [2](#) respectively. However, it is of interest to consider more general inequalities of Janous-type and of Tsintsifas-type which may hold under the weak condition $A + B + C = \theta$ where $0 < \theta \leq \pi$. We are inspired by the following parameterized Jordan-type inequality presented in our recent paper [22] to pursue the proposed results.

Theorem B (See Wu and Debnath [22]). Let $0 < x \leq \theta \leq \frac{\pi}{2}$. Then

$$\frac{\sin \theta}{\theta} - \frac{\theta - \sin \theta}{\theta^2} (x - \theta) \leq \frac{\sin x}{x} \leq \frac{\sin \theta}{\theta} + \frac{\theta \cos \theta - \sin \theta}{\theta^2} (x - \theta). \quad (4)$$

It is important to note that the inequality (4) cannot be applied in the improvement of inequalities (2) and (3) because the condition of parameter is $0 < \theta \leq \pi/2$ rather than $0 < \theta \leq \pi$ which is what we would need for the inequality to be applied. Therefore, a further study on the Jordan-type inequality (4) is needed.

In this work, we shall generalize the inequality (4). Under certain weak constraint conditions, we prove that inequality (4) is still valid. In Section 3, the result will be used to generalize Janous's inequality (2) and improve Tsintsifas's inequality (3).

2. Generalization of Jordan-type inequality

In the generalization of Jordan-type inequality (4), we relax the constraint conditions of parameter in (4), the main result is the following theorem:

Theorem 1. If $0 < x \leq \theta \leq \pi$, then

$$\frac{\sin x}{x} \geq \frac{\sin \theta}{\theta} - \frac{\theta - \sin \theta}{\theta^2} (x - \theta). \quad (5)$$

If $0 < x \leq \pi$ and $0 < \theta \leq \frac{\pi}{2}$, then

$$\frac{\sin x}{x} \leq \frac{\sin \theta}{\theta} + \frac{\theta \cos \theta - \sin \theta}{\theta^2} (x - \theta). \quad (6)$$

Furthermore, the equalities hold in (5) or (6) if and only if $x = \theta$.

Proof. We introduce a function

$$f : (0, \pi] \longrightarrow \mathbb{R}$$

by

$$f(x) = \frac{1}{x} \left(\frac{\sin x}{x} - 1 \right).$$

Differentiating $f(x)$ with respect to x gives

$$f'(x) = \frac{x \cos x + x - 2 \sin x}{x^3}.$$

By means of a simple calculation and the well-known inequality $\tan \alpha > \alpha$ ($0 < \alpha < \pi/2$), we find that for $0 < x < \pi$,

$$f'(x) = \frac{x \cos x + x - 2 \sin x}{x^3} = \frac{4 \cos^2 \frac{x}{2} \left(\frac{x}{2} - \tan \frac{x}{2} \right)}{x^3} < 0.$$

This means that $f(x)$ is strictly decreasing on $(0, \pi)$. Thus, we deduce that for $0 < x < \theta \leq \pi$,

$$\frac{1}{x} \left(\frac{\sin x}{x} - 1 \right) > \frac{1}{\theta} \left(\frac{\sin \theta}{\theta} - 1 \right),$$

which can be simplified to the desired inequality:

$$\frac{\sin x}{x} > \frac{\sin \theta}{\theta} - \frac{\theta - \sin \theta}{\theta^2} (x - \theta). \quad (7)$$

In addition, it is easy to see that the equality occurs in (5) if and only if $x = \theta$. The inequality (5) is thus proved.

We next define a function

$$g : (0, \pi] \longrightarrow \mathbb{R}$$

by

$$g(x) = \frac{\sin x}{x} - \frac{\theta \cos \theta - \sin \theta}{\theta^2} (x - \theta) - \frac{\sin \theta}{\theta}.$$

Differentiating $g(x)$ with respect to x gives

$$g'(x) = \frac{x \cos x - \sin x}{x^2} - \frac{\theta \cos \theta - \sin \theta}{\theta^2},$$

$$g''(x) = \frac{2 \sin x - 2x \cos x - x^2 \sin x}{x^3}.$$

Let $\varphi(x) = 2 \sin x - 2x \cos x - x^2 \sin x$, then $\varphi'(x) = -x^2 \cos x$. In view of the fact that $\varphi'(x) < 0$ for $0 < x < \frac{\pi}{2}$ and $\varphi'(x) > 0$ for $\frac{\pi}{2} < x < \pi$, we conclude that $\varphi(x)$ is strictly decreasing on $(0, \frac{\pi}{2})$ and strictly increasing on $(\frac{\pi}{2}, \pi)$.

Further, we deduce from the observation $\varphi(\frac{\pi}{2}) < 0$ and $\varphi(\pi) > 0$ that there is a point $x_1 \in (\frac{\pi}{2}, \pi)$ such that $\varphi(x) < 0$ for $0 < x < x_1$ and $\varphi(x) > 0$ for $x_1 < x < \pi$. We then conclude that $g'(x)$ is strictly decreasing on $(0, x_1)$ and strictly increasing on (x_1, π) .

In addition, it follows from straightforward computation that

$$g'(0+) = \frac{\cos \theta (\tan \theta - \theta)}{\theta^2} > 0, \quad g'(\theta) = 0 \quad (0 < \theta \leq \pi/2 < x_1), \quad (8)$$

which implies that

$$g'(x_1) < g'\left(\frac{\pi}{2}\right) \leq 0. \quad (9)$$

To prove the required inequality (6), we consider the following two cases.

Case (I): When $g'(\pi) \leq 0$.

By relations (8), (9), $g'(\pi) \leq 0$ and the monotonicity of $g'(x)$, it follows that $g'(x) > 0$ for $0 < x < \theta$ and $g'(x) < 0$ for $\theta < x < \pi$. We hence conclude that $g(x)$ is strictly increasing on $(0, \theta)$ and strictly decreasing on (θ, π) .

Therefore, we have

$$g(x) \leq g(\theta) = 0 \quad \text{for } 0 < x \leq \pi, \quad (10)$$

which leads us to the desired inequality (6).

Case (II): When $g'(\pi) > 0$.

Since $g'(x_1) < 0$, $g'(\pi) > 0$ and $g'(x)$ is strictly increasing on (x_1, π) , there exists a point $x_2 \in (x_1, \pi)$ such that $g'(x) < 0$ for $x_1 < x < x_2$ and $g'(x) > 0$ for $x_2 < x < \pi$. On the other hand, we observe via relations (8) and (9) that $g'(x) > 0$ for $0 < x < \theta$ and $g'(x) < 0$ for $\theta < x < x_1$.

Consequently, $g(x)$ is strictly increasing on $(0, \theta)$, strictly decreasing on (θ, x_2) and strictly increasing on (x_2, π) .

Now, from $g(\theta) = 0$ and the following inequality holds:

$$\begin{aligned} g(\pi) &= -\frac{\theta \cos \theta - \sin \theta}{\theta^2} (\pi - \theta) - \frac{\sin \theta}{\theta} \\ &= \frac{\pi}{2} \left(-\frac{4}{\pi^2} - \frac{\theta \cos \theta - \sin \theta}{\theta^2} \right) + \left(\frac{2}{\pi} - \frac{\theta \cos \theta - \sin \theta}{\theta^2} \left(\frac{\pi}{2} - \theta \right) - \frac{\sin \theta}{\theta} \right) \\ &= \frac{\pi}{2} g'\left(\frac{\pi}{2}\right) + g\left(\frac{\pi}{2}\right) \leq \frac{\pi}{2} g'(\theta) + g(\theta) = 0. \end{aligned}$$

Finally, we deduce that

$$g(x) \leq 0 \quad \text{for } 0 < x \leq \pi, \quad (11)$$

which implies the desired inequality (6). This completes the proof of Theorem 1. \square

Remark 1. It is obvious that the Jordan-type inequality (4) will follow as a special case of Theorem 1 when $0 < x \leq \theta \leq \pi/2$.

3. Some applications

As applications of Theorem 1, Janous's inequality (2) and Tsintsifas's inequality (3) mentioned in Section 1 will be generalized and improved in this section.

Theorem 2. Let $x_i > 0$ ($i = 1, 2, \dots, n$, $n \geq 2$), $x_1 + x_2 + \dots + x_n = \theta$ and $0 < \theta \leq \pi$. Then

$$\frac{\sin \theta}{\theta} + n - 1 < \sum_{i=1}^n \frac{\sin x_i}{x_i} \leq \frac{n^2}{\theta} \sin \frac{\theta}{n}, \quad (12)$$

with equality holding if and only if $x_1 = x_2 = \dots = x_n = \theta/n$.

Proof. Since $0 < x_i < \theta \leq \pi$ ($i = 1, 2, \dots, n$), we obtain via inequality (5) that

$$\frac{\sin x_i}{x_i} > \frac{\sin \theta}{\theta} - \frac{\theta - \sin \theta}{\theta^2} (x_i - \theta) \quad (i = 1, 2, \dots, n). \quad (13)$$

Taking the sum for all inequalities of (13) yields

$$\begin{aligned} \sum_{i=1}^n \frac{\sin x_i}{x_i} &> \frac{n \sin \theta}{\theta} - \frac{\theta - \sin \theta}{\theta^2} \sum_{i=1}^n (x_i - \theta) \\ &= \frac{\sin \theta}{\theta} + n - 1. \end{aligned} \quad (14)$$

The hypothesis of Theorem 2 shows that

$$0 < x_i \leq \pi, \quad 0 < \theta/n \leq \pi/2, \quad i = 1, 2, \dots, n, \quad n \geq 2.$$

Hence, from inequality (6) we deduce that

$$\frac{\sin x_i}{x_i} \leq \frac{\sin \frac{\theta}{n}}{\frac{\theta}{n}} + \frac{\frac{\theta}{n} \cos \frac{\theta}{n} - \sin \frac{\theta}{n}}{\left(\frac{\theta}{n}\right)^2} \left(x_i - \frac{\theta}{n}\right) \quad (i = 1, 2, \dots, n), \quad (15)$$

where the equality holds if and only if $x = \theta/n$.

Taking the sum for all inequalities of (15), we get

$$\begin{aligned} \sum_{i=1}^n \frac{\sin x_i}{x_i} &\leq \frac{n \sin \frac{\theta}{n}}{\frac{\theta}{n}} + \frac{\frac{\theta}{n} \cos \frac{\theta}{n} - \sin \frac{\theta}{n}}{\left(\frac{\theta}{n}\right)^2} \sum_{i=1}^n \left(x_i - \frac{\theta}{n}\right) \\ &= \frac{n^2}{\theta} \sin \frac{\theta}{n}. \end{aligned}$$

This completes the proof of Theorem 2. \square

Setting $n = 3$, $x_1 = A$, $x_2 = B$, $x_3 = C$ in Theorem 2, a generalization of Janous's inequality (2) is derived as follows:

Corollary 1. Let $A > 0$, $B > 0$, $C > 0$, $A + B + C = \theta$ and $0 < \theta \leq \pi$. Then

$$\frac{\sin \theta}{\theta} + 2 < \frac{\sin A}{A} + \frac{\sin B}{B} + \frac{\sin C}{C} \leq \frac{9}{\theta} \sin \frac{\theta}{3}. \quad (16)$$

It is important to point out that Corollary 1 with the special case $\theta = \pi$ yields Janous's inequality (2).

Theorem 3. Let $x_i > 0$ ($i = 1, 2, \dots, n$, $n \geq 2$), $x_1 + x_2 + \dots + x_n = \theta$ and $0 < \theta \leq \pi$. Then

$$\sum_{i=1}^n \frac{\sin x_i}{\theta - x_i} > \frac{1}{n-1} \left(\frac{\sin \theta}{\theta} \right) + 1. \quad (17)$$

Proof. Note that $0 < x_i < \theta \leq \pi$ ($i = 1, 2, \dots, n$), then it follows from inequality (5) that

$$\begin{aligned} \frac{\sin x_i}{\theta - x_i} &> \left(\frac{\sin \theta}{\theta} \right) \frac{x_i}{\theta - x_i} + \left(\frac{\theta - \sin \theta}{\theta^2} \right) x_i \\ &= \frac{\sin \theta}{\theta - x_i} + \left(\frac{\theta - \sin \theta}{\theta^2} \right) x_i - \frac{\sin \theta}{\theta} \quad (i = 1, 2, \dots, n). \end{aligned} \quad (18)$$

Summing up (18) for $i = 1, 2, \dots, n$ yields

$$\begin{aligned} \sum_{i=1}^n \frac{\sin x_i}{\theta - x_i} &> \sin \theta \left(\sum_{i=1}^n \frac{1}{\theta - x_i} \right) + \left(\frac{\theta - \sin \theta}{\theta^2} \right) \left(\sum_{i=1}^n x_i \right) - \frac{n \sin \theta}{\theta} \\ &= \sin \theta \left(\sum_{i=1}^n \frac{1}{\theta - x_i} \right) - \frac{(n+1) \sin \theta}{\theta} + 1. \end{aligned} \quad (19)$$

Applying the Cauchy–Schwarz inequality, we obtain

$$\sum_{i=1}^n \frac{1}{\theta - x_i} \geq n^2 \left(\sum_{i=1}^n (\theta - x_i) \right)^{-1} = \frac{n^2}{(n-1)\theta}. \quad (20)$$

Substituting (20) into (19) leads to inequality (17). The proof of Theorem 3 is complete. \square

Setting $n = 3$, $x_1 = A$, $x_2 = B$, $x_3 = C$ in Theorem 3, we get the generalized version of Tsintsifas's inequality (3):

Corollary 2. Let $A > 0$, $B > 0$, $C > 0$, $A + B + C = \theta$ and $0 < \theta \leq \pi$. Then

$$\frac{\sin A}{\theta - A} + \frac{\sin B}{\theta - B} + \frac{\sin C}{\theta - C} > \frac{\sin \theta}{2\theta} + 1. \quad (21)$$

If we put $\theta = \pi$ in inequality (21), we obtain the following result:

$$\frac{\sin A}{\pi - A} + \frac{\sin B}{\pi - B} + \frac{\sin C}{\pi - C} > 1. \quad (22)$$

Inequality (22) partially improves Tsintsifas's inequality (3), since the lower bound of (22) is sharper than that of (3).

Theorem 4. Let $x_i > 0$ ($i = 1, 2, \dots, n$, $n \geq 3$), $x_1 + x_2 + \dots + x_n = \theta$ and $0 < \theta \leq \pi$. Then

$$(n-1) \left(\frac{\sin \theta}{\theta} \right) + 1 < \sum_{i=1}^n \frac{\sin(\theta - x_i)}{\theta - x_i} < (n^2 - 3n + 1) \cos \frac{\theta}{n-1} - \frac{(n-1)(n^2 - 4n + 1)}{\theta} \sin \frac{\theta}{n-1}. \quad (23)$$

Proof. The hypothesis of Theorem 4 implies that

$$0 < \theta - x_i < \theta \leq \pi, \quad 0 < \theta/(n-1) \leq \pi/2, \quad i = 1, 2, \dots, n, \quad n \geq 3.$$

Thus, using inequalities (5) and (6), we obtain that for $i = 1, 2, \dots, n$,

$$\begin{aligned} \frac{\sin \theta}{\theta} - \frac{\theta - \sin \theta}{\theta^2} (\theta - x_i - \theta) &< \frac{\sin(\theta - x_i)}{\theta - x_i} \\ &< \frac{\sin \frac{\theta}{n-1}}{\frac{\theta}{n-1}} + \frac{\frac{\theta}{n-1} \cos \frac{\theta}{n-1} - \sin \frac{\theta}{n-1}}{\left(\frac{\theta}{n-1}\right)^2} \left(\theta - x_i - \frac{\theta}{n-1} \right). \end{aligned} \quad (24)$$

Summing up (24) for $i = 1, 2, \dots, n$ yields

$$\begin{aligned} \frac{n \sin \theta}{\theta} - \frac{\theta - \sin \theta}{\theta^2} \sum_{i=1}^n (\theta - x_i - \theta) &< \sum_{i=1}^n \frac{\sin(\theta - x_i)}{\theta - x_i} \\ &< \frac{n \sin \frac{\theta}{n-1}}{\frac{\theta}{n-1}} + \frac{\frac{\theta}{n-1} \cos \frac{\theta}{n-1} - \sin \frac{\theta}{n-1}}{\left(\frac{\theta}{n-1}\right)^2} \sum_{i=1}^n \left(\theta - x_i - \frac{\theta}{n-1} \right). \end{aligned} \quad (25)$$

A simple computation reduces inequality (25) to inequality (23). The proof of Theorem 4 is complete. \square

Setting $n = 3$, $x_1 = A$, $x_2 = B$, $x_3 = C$ in Theorem 4 yields the following generalized and sharpened version of Tsintsifas's inequality (3).

Corollary 3. Let $A > 0$, $B > 0$, $C > 0$, $A + B + C = \theta$ and $0 < \theta \leq \pi$. Then

$$\frac{2 \sin \theta}{\theta} + 1 < \frac{\sin(\theta - A)}{\theta - A} + \frac{\sin(\theta - B)}{\theta - B} + \frac{\sin(\theta - C)}{\theta - C} < \cos \frac{\theta}{2} + \frac{4}{\theta} \sin \frac{\theta}{2}. \quad (26)$$

In particular, putting $\theta = \pi$ in Corollary 3, we obtain:

Corollary 4. Let $A > 0$, $B > 0$, $C > 0$ and $A + B + C = \pi$. Then

$$1 < \frac{\sin A}{\pi - A} + \frac{\sin B}{\pi - B} + \frac{\sin C}{\pi - C} < \frac{4}{\pi}. \quad (27)$$

Obviously, inequality (27) is stronger than Tsintsifas's inequality (3), and also, it is more general than Tsintsifas's result.

Acknowledgment

The research was supported, in part, by the Natural Science Foundation of Fujian province of China under grant No. S0650003 and, in part, by the program for new century excellent talents in the newly-built universities of Fujian province of China.

References

- [1] D.S. Mitrinović, P.M. Vasić, *Analytic Inequalities*, Springer-Verlag, New York, 1970.
- [2] S. Wu, On generalizations and refinements of Jordan type inequality, *Octagon Math. Mag.* 12 (1) (2004) 267–272.
- [3] S. Wu, L. Debnath, A new generalized and sharp version of Jordan's inequality and its applications to the improvement of the Yang Le inequality, II, *Appl. Math. Lett.* 20 (5) (2007) 532–538.
- [4] S. Wu, Sharpness and generalization of Jordan's inequality and its application, *Taiwanese J. Math.* 12 (2) (2008) 325–336.
- [5] L. Zhu, Sharpening Jordan's inequality and Yang Le inequality, *Appl. Math. Lett.* 19 (3) (2006) 240–243.
- [6] L. Zhu, Sharpening of Jordan's inequalities and its applications, *Math. Inequal. Appl.* 9 (1) (2006) 103–106.
- [7] L. Zhu, Sharpening Jordan's inequality and Yang Le inequality, II, *Appl. Math. Lett.* 19 (9) (2006) 990–994.
- [8] A.Y. Özban, A new refined form of Jordan's inequality and its applications, *Appl. Math. Lett.* 19 (2006) 155–160.
- [9] L. Debnath, C.J. Zhao, New strengthened Jordan's inequality and its applications, *Appl. Math. Lett.* 16 (4) (2003) 557–560.
- [10] C.J. Zhao, L. Debnath, On Generalizations of L. Yang's inequality, *J. Inequal. Pure Appl. Math.* 3 (4) (2002) Article 56, pp. 1–5 (electronic).
- [11] W.D. Jiang, H. Yun, Sharpening of Jordan's inequality and its applications, *J. Inequal. Pure Appl. Math.* 7 (3) (2006) Article 102, pp. 1–8 (electronic).
- [12] X. Zhang, G. Wang, Y. Chu, Extensions and sharpening of Jordan's inequality and Kober's inequality, *J. Inequal. Pure Appl. Math.* 7 (2) (2006) Article 63, pp. 1–7 (electronic).
- [13] F. Yuefeng, Jordan's inequality, *Math. Mag.* 69 (1996) 126–127.
- [14] A.McD. Mercer, U. Abel, D. Caccia, A sharpening of Jordan's inequality, *Amer. Math. Monthly* 93 (1986) 568–569.
- [15] S. Wu, L. Debnath, A new generalized and sharp version of Jordan's inequality and its applications to the improvement of the Yang Le inequality, *Appl. Math. Lett.* 19 (12) (2006) 1378–1384.
- [16] S. Wu, H.M. Srivastava, A weighted and exponential generalization of Wilker's inequality and its applications, *Integral Transforms. Spec. Funct.* 18 (2007) 529–535.
- [17] S. Wu, H.M. Srivastava, L. Debnath, Some refined families of Jordan type inequalities and their applications, *Integral Transforms. Spec. Funct.* 19 (2008) 183–193.
- [18] W. Janous, Problem 1216, *Crux Math.* 13 (1987) 53.
- [19] G. Tsintsifas, Problem 1365, *Crux Math.* 14 (1988) 202.
- [20] M.S. Klamkin, Solution of problem 1216, *Crux Math.* 14 (1988) 120–122.
- [21] V.N. Murty, G.P. Henderson, Solution of problem 1365, *Crux Math.* 15 (1989) 251–256.
- [22] S. Wu, L. Debnath, Jordan-type inequalities for differentiable functions and their applications, *Appl. Math. Lett.* 21 (8) (2008) 803–809.